



Patient Safety Standing Committee Web Meeting

Andrew Lyzenga
Jesse Pines
Kathryn Goodwin
Hiral Dudhwala
Desmirra Quinnonez

September 12, 2018

Agenda for Today's Meeting

- Welcome/Roll Call
- Guidance on Harmonization of Medication Reconciliation Measures
- Additional Patient Safety Topics for Harmonization
- NQF Member and Public Comment
- Next Steps
- Adjourn

Patient Safety Project Team



Andrew Lyzenga
Senior Director



Jesse Pines
Consultant



Kathryn Goodwin
Senior Project Manager



Hiral Dudhwala
Project Manager



Desmirra Quinnonez
Project Analyst

Patient Safety Standing Committee

- Ed Septimus, MD (Co-Chair)
- Iona Thraen, PhD, ACSW (Co-Chair)
- Jason Adelman, MD, MS
- Charlotte Alexander, MD
- Kimberly Applegate, MD, MS, FACR
- Laura Ardizzone, BSN, MS, DNP, CRNA
- Richard Brill, MD, FAAP, FCCM
- Curtis Collins, PharmD, MS
- Christopher Cook, PharmD, PhD
- Melissa Danforth, BA
- Theresa Edelstein, MPH, LNHA
- Lillee Gelinas, MSN, RN, FAAN
- John James, PhD
- Stephen Lawless, MD, MBA, FAAP, FCCM
- Lisa McGiffert
- Susan Moffatt-Bruce, MD, PhD
- Patricia Quigley, PhD, MPH, ARNP, CRRN, FAAN, FAANP
- Michelle Schreiber, MD
- Leslie Schultz, PhD, RN, NEA-BC, CPHQ
- Lynda Smirz, MD, MBA
- Tracy Wang, MPH
- Kendall Webb, MD, FACEP
- Albert Wu, MD, MPH, FACP
- Donald Yealy, MD, FACEP
- Yanling Yu, PhD

Patient Safety Standing Committee Expert Reviewers

* New Expert Reviewer

- Jamie Roney, DNP, RN-BC, CCRN-K
 - *(Infectious Disease)*
- Pranavi Sreeramoju, MD, MPH, CMQ, FSHEA, FIDSA
 - *(Infectious Disease)*
- Bruno Digiovine, MD
 - *(Pulmonary)*
- Edgar Jimenez, MD, FCCM
 - *(Pulmonary)*
- David Stockwell, MD, MBA
 - *(Pulmonary)*
- Emily Aaronson, MD*
 - *(Infectious Disease)*

Guidance on Harmonization of Medication Reconciliation Measures

Context

- Fall 2017 Behavioral Health SC discussion about medication reconciliation
 - Desire for greater alignment in measure specifications
- April 2018 CSAC meeting
 - *Medication reconciliation is a general topic*
 - *Which is best?*
 - » Narrowly focused measures (e.g., med rec for a specific patient group) **OR**
 - » Broader measure that includes most patients

Good opportunity to talk about our processes for related and competing measures more generally

Definitions

At the **conceptual** level:

Competing Measures	Related Measures
Same measure focus AND Same target population	Same measure focus OR Same target population



Harmonize if possible
(align specifications)

Exemplar: Flu shot measures

- 2008 Steering Committee identified standard measure specifications
 - *Who is included in/excluded from the target denominator population*
 - *Who is included in the numerator population*
 - *Time windows for measurement and vaccinations*
 - *Exclusions*
- 2012 Population Health Steering Committee strongly recommended the development of a universal influenza immunization measure
- 2017 Health and Well-Being Standing Committee
 - *Evaluated and endorsed eight flu measures*
 - » Most harmonized to NQF's standardized specifications
 - » SC reiterated the need for a single, standardized measure

Review of Evidence in NQF-Endorsed Medication Reconciliation Measures

- No clinical guidelines cited
- Several individual studies that have demonstrated a decrease in medication errors when medication reconciliation is implemented
- Several articles on adverse drug events in the various care settings
- 2012 systematic review on Hospital-Based Medication Reconciliation Practices*
 - *Reduction of medication discrepancies and ADEs and successful interventions include: intensive pharmacy staff involvement and targeting intervention to “high risk” patient population.*

*Mueller SK, Sponsler KC, Kripalani S, Schnipper JL. Hospital-Based Medication Reconciliation Practices: A Systematic. *Arch Intern Med.* Jun 25 2012:1-13.

Review of Evidence in NQF-Endorsed Medication Reconciliation Measures

- Joint Commission's Ambulatory and Hospital Patient Safety Goal recommendations
 - *NPSG.03.06.01: "Maintain and communicate accurate patient medication information. The types of information that clinicians use to reconcile medications include (among others) medication name, dose, frequency, route, and purpose. Organizations should identify the information that needs to be collected to reconcile current and newly ordered medications and to safely prescribe medications in the future." (Ambulatory)*
 - *"Obtain information on the medications the patient is currently taking when he or she is admitted to the hospital or is seen in an outpatient setting. This information is documented in a list or other format that is useful to those who manage medications. (Hospital)*

Related Medication Reconciliation Measures

	0097: MedRec Post-Discharge	0419e: Documentation of Current Medications in the Medical Record	0553: Care for Older Adults (COA) – Medication Review	2456: MedRec: Number of Unintentional Medication Discrepancies per Patient	3317: MedRec on Admission	2988: MedRec for Patients Receiving Care at Dialysis Facilities
Steward	NCQA	CMS	NCQA	Brigham and Women’s Hospital	CMS / HSAG	Kidney Quality Care Alliance
Numerator	Medication reconciliation conducted by a prescribing practitioner, clinical pharmacist or registered nurse on or within 30 days of discharge. Medication reconciliation is defined as a type of review in which the discharge medications are reconciled with the most recent medication list in the outpatient medical record.	The Numerator statement for 2017 Claims/Registry Specification: Eligible clinician attests to documenting, updating, or reviewing a patient’s current medications using all immediate resources available on the date of the encounter. This list must include ALL prescriptions, over-the counters, herbals, vitamin/mineral/dietary (nutritional) supplements AND must contain the medications’ name, dosages, frequency, and route of administration. 2018 eMeasure Specification: Eligible professional or eligible clinician attests to documenting, updating or reviewing the patient’s current medications using all immediate resources available on the date of the encounter. This list must include ALL known prescriptions, over-the-counters, herbals and vitamin/mineral/dietary (nutritional) supplements AND must contain the medications’ name, dosages, frequency and route of administration.	At least one medication review conducted by a prescribing practitioner or clinical pharmacist during the measurement year and the presence of a medication list in the medical record	For each sampled inpatient in the denominator, the total number of unintentional medication discrepancies in admission orders plus the total number of unintentional medication discrepancies in discharge orders.	Number of patients for whom a designated Prior to Admission (PTA) medication list was generated by referencing one or more external sources of medications and for which all PTA medications have a documented reconciliation action by the end of Day 2 of the hospitalization when the admission date is Day 0.	Number of patient-months for which medication reconciliation was performed and documented by an eligible professional during the reporting period. The medication reconciliation MUST: <ul style="list-style-type: none"> • Include the name or other unique identifier of the eligible professional; AND • Include the date of the reconciliation; AND • Address ALL known home medications (prescriptions, over-the-counters, herbals, vitamin/mineral/dietary (nutritional) supplements, and medical marijuana); AND • Address for EACH home medication: Medication name(1), indication(2), dosage(2), frequency(2), route of administration(2), start and end date (if applicable)(2), discontinuation date (if applicable)(2), reason medication was stopped or discontinued (if applicable)(2), and identification of individual who authorized stoppage or discontinuation of medication (if applicable)(2); AND • List any allergies, intolerances, or adverse drug events experienced by the patient. <ol style="list-style-type: none"> 1. For patients in a clinical trial, it is acknowledged that it may be unknown as to whether the patient is receiving the therapeutic agent or a placebo. 2. “Unknown” is an acceptable response for this field.

	0097: MedRec Post-Discharge	0419e: Documentation of Current Medications in the Medical Record	0553: Care for Older Adults (COA) – Medication Review	2456: MedRec: Number of Unintentional Medication Discrepancies per Patient	3317: MedRec on Admission	2988: MedRec for Patients Receiving Care at Dialysis Facilities
Denominator	All discharges from an in-patient setting for patients who are 18 years and older.	The 2017 Claims and Registry denominator statement is as follows: “All visits for patients aged 18 years and older.” The 2018 eMeasure denominator statement is as follows: “All visits occurring during the 12 month reporting measurement period for patients aged 18 years and older.”	All patients 66 and older as of the end (e.g., December 31) of the measurement year.	The patient denominator includes a random sample of all potential adults admitted to the hospital. Our recommendation is that 25 patients are sampled per month, or approximately 1 patient per weekday. So, for example, if among those 25 patients, 75 unintentional discrepancies are identified, the measure outcome would be 3 discrepancies per patient for that hospital for that month.	All patients admitted to an inpatient facility from home or a non-acute setting.	Total number of patient-months for all patients permanently assigned to a dialysis facility during the reporting period.
Exclusions	The following exclusions are applicable to the Health Plan Level measure. - Exclude both the initial discharge and the readmission/direct transfer discharge if the readmission/direct transfer discharge occurs after December 1 of the measurement year. - If the discharge is followed by a readmission or direct transfer to an acute or non-acute facility within the 30-day follow-up period, count only the readmission discharge or the discharge from the facility to which the patient was transferred.	The 2017 Claims and Registry version provides the following as a denominator exception: A patient is not eligible if the following reason is documented: Patient is in an urgent or emergent medical situation where time is of the essence and to delay treatment would jeopardize the patient’s health status. The 2018 eMeasure includes the following denominator exception: Medical Reason: Patient is in an urgent or emergent medical situation where time is of the essence and to delay treatment would jeopardize the patient’s health status.	Exclude members who use hospice services or elect to use a hospice benefit any time during the measurement year, regardless of when the services began.	Patients that are discharged or expire before a gold standard medication list can be obtained.	The measure applies two exclusion criteria to ensure that it is feasible to complete the medication reconciliation process on admission to the IPF: 1. Patients transferred from an acute care setting 2. Patient admissions with a length of stay less than or equal to 2 days	In-center patients who receive <7 hemodialysis treatments in the facility during the reporting month.

	0097: MedRec Post-Discharge	0419e: Documentation of Current Medications in the Medical Record	0553: Care for Older Adults (COA) – Medication Review	2456: MedRec: Number of Unintentional Medication Discrepancies per Patient	3317: MedRec on Admission	2988: MedRec for Patients Receiving Care at Dialysis Facilities
Measure Focus	Reconciliation of discharge medication list with current outpatient medical record medication list	Eligible clinician attests to documenting a list of current medications using all immediate resources available on the date of the encounter	Medication review of all a patient’s medications, including prescription medications, OTC medications by a prescribing practitioner or clinical pharmacist	Total number of unintentional medication discrepancies in admission orders + total number of unintentional medication discrepancies in discharge orders	Reconciliation of Prior to Admission medication list (referencing external sources) by end of Day 2 of hospitalization.	Patients receive medication reconciliation upon visit to dialysis facility.
Population	Patients ages 18 +	Patients ages 18 +	Patients ages 66 +	Random sample of adults admitted to the hospital	All inpatient psychiatric admissions	Dialysis patients
Data Source	Claims, Electronic Health Records, Paper Medical Records	Claims, Electronic Health Records, Registry Data	Claims, Electronic Health Records, Paper Medical Records	Electronic Health Data, Electronic Health Records, Instrument-Based Data, Other, Paper Medical Records	Paper Medical Records	Electronic Health Records, Other
Level of Analysis	Clinician: individual Clinician: group Health Plan Integrated Delivery System	Clinician: individual Clinician: group	Health Plan Integrated Delivery System	Facility	Facility	Facility
Setting	Outpatient	Outpatient	Inpatient/Hospital, Outpatient Services, Post-Acute Care	Hospital	Inpatient/Hospital	Post-Acute Care

Medication Reconciliation: Issues to Consider

- What would be included in standardized specifications?
 - *What would be reconciled?*
 - » All prescriptions, OTCs, herbals, vitamins, etc.
 - » Name, dosages, frequency, route
 - *How often does it need to be done?*
 - *Who would do it? (e.g., pharmacist, MDs, etc.)*
 - *Who needs it done? (e.g., all pts? Stratify for certain groups?)*
 - *What would trigger it? (e.g., “visit”, phone refill, etc.)*
 - Where should it be done?
 - What terminology coding should be used (SNOMED, LOINC, RXNorm)?
- Is there any evidence to inform the above?
 - *Does it differ across settings, patient populations, or conditions?*
- What might differ depending on care setting, data source, level of analysis?

Medication Reconciliation: Issues to Consider

- Where should medication reconciliation measurement go in the future?
 - *Outcome measures, adverse events?*
 - *What is the ideal future state of measurement with respect to medication reconciliation?*
 - *What research might be needed to support future quality measurement in this area?*
- How can we best harmonize existing med rec measures?

Additional Patient Safety Topics for Harmonization

Public Comment

Next Steps

Activities and Timeline

Fall 2017 Cycle

Process Step	Timeline
Final Technical Report	July 23, 2018 (POSTED)

Spring 2018 Cycle

Process Step	Timeline
Committee Web Meeting	Wednesday, September 12, 2018 1-3 PM EST (TODAY)

Activities and Timeline

Fall 2018 Cycle

Process Step	Timeline
Intent to Submit Deadline (Measure Steward/Developers)	Wednesday, August 1, 2018
Measure Submission Deadline (Measure Steward/Developers)	Thursday, November 1, 2018

Fall 2018 Measures

- 7 Maintenance Measures:
 - **NQF #0553** *Care for Older Adults (COA) – Medication Review*
 - **NQF #0555** *INR Monitoring for Individuals on Warfarin*
 - **NQF #0753** *American College of Surgeons – Centers for Disease Control and Prevention (ACS-CDC) Harmonized Procedure Specific Surgical Site Infection (SSI) Outcome Measure*
 - **NQF #1716** *National Healthcare Safety Network (NHSN) Facility-wide Inpatient Hospital-onset Methicillin-resistant Staphylococcus aureus (MRSA) Bacteremia Outcome Measure*
 - **NQF #1717** *National Healthcare Safety Network (NHSN) Facility-wide Inpatient Hospital-onset Clostridium difficile Infection (CDI) Outcome Measure*
 - **NQF #2456** *Medication Reconciliation: Number of Unintentional Medication Discrepancies per Patient*
 - **NQF #3450** *Practice Environment Scale - Nursing Work Index (PES-NWI) (composite and five subscales)*

THANK YOU

Adjourn